ANTIBODIES ONLINE

Datasheet for ABIN928550 anti-SERTAD2 antibody (N-Term)

Image



Overview


1

| Quantity: | 100 µL |
|----------------------|---|
| Target: | SERTAD2 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse, Rat, Zebrafish (Danio rerio), Chicken, Dog, Xenopus laevis |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SERTAD2 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | SERTAD2 antibody was raised in rabbit using the N terminal of SERTAD2 as the immunogen |
| Cross-Reactivity: | Mouse (Murine), Rat (Rattus), Dog (Canine), Chicken, Frog, Zebrafish (Brachydanio rerio) |
| Purification: | Purified |
| Target Details | |
| Target: | SERTAD2 |
| Alternative Name: | SERTAD2 (SERTAD2 Products) |
| Background: | SERTAD2 acts at E2F-responsive promoters to integrate signals provided by PHD- and/or |
| | bromodomain-containing transcription factors. Synonyms: Polyclonal SERTAD2 antibody, Anti |
| | |
| | SERTAD2 antibody, SERTA domain containing 2 antibody, MGC126688 antibody, MGC126690 |

International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN928550 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

| Application Details | |
|---------------------|--|
| Application Notes: | WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator. |
| Comment: | SERTAD2 Blocking Peptide, catalog no. 33R-5336, is also available for use as a blocking control in assays to test for specificity of this SERTAD2 antibody |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Concentration: | Lot specific |
| Buffer: | Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. |

Images



Western Blotting

Image 1. SERTAD2 antibody used at 0.2-1 ug/ml to detect target protein.