## ANTIBODIES ONLINE

## Datasheet for ABIN928567 anti-TP53I13 antibody (C-Term)

Image



Overview


1

01011010	
Quantity:	100 µL
Target:	TP53I13
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TP53I13 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	TP53 I13 antibody was raised in rabbit using the C terminal of TP53 13 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Dog (Canine)
Purification:	Purified
Target Details	
Target:	TP53I13
Alternative Name:	TP53I13 (TP53I13 Products)
Background:	TP53I13 may act as a tumor suppressor. It inhibits tumor cell growth, when overexpressed.
	TP53I13 was up-regulated by genotoxic stresses of adriamycin and/or ultraviolet beam
	irradiation in a p53-dependent manner. Synonyms: Polyclonal TP53I13 antibody, Anti-TP53I13
	antibody, tumor protein p53 inducible protein 13 antibody, DSCP1 antibody.
Order at www :	antibodies-online.com   www.antikoerper-online.de   www.anticorps-enligne.fr   www.antibodies-online.cn

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN928567 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Application Details	
Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	TP53I13 Blocking Peptide, catalog no. 33R-10291, is also available for use as a blocking control in assays to test for specificity of this TP53I13 antibody
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Format: Concentration:	Lyophilized Lot specific
Concentration:	Lot specific Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in
Concentration: Buffer:	Lot specific Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

## Images



## Western Blotting

**Image 1.** TP53I13 antibody used at 0.2-1 ug/ml to detect target protein.