

Datasheet for ABIN928665

anti-FBXO32 antibody (N-Term)





Overview

| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | FBXO32 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FBXO32 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | FBXO32 antibody was raised in rabbit using the N terminal of FBXO32 as the immunogen |
| Purification: | Purified |
| Target Details | |
| Target: | FBXO32 |
| Alternative Name: | FBXO32 (FBXO32 Products) |
| Background: | This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs, which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 |
| | domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein- |

protein interaction modules or no recognizable motifs. Synonyms: Polyclonal FBXO32 antibody, Anti-FBXO32 antibody, F-box protein 32 antibody, FLJ32424 antibody, Fbx32 antibody, MAFbx antibody, MGC33610 antibody.


Application Details

| Application Notes: | WB: 0.2-1 μg/mL |
|--------------------|--|
| | Optimal conditions should be determined by the investigator. |
| Comment: | FBXO32 Blocking Peptide, catalog no. 33R-7699, is also available for use as a blocking control in assays to test for specificity of this FBXO32 antibody |
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized |
|------------------|---|
| Concentration: | Lot specific |
| Buffer: | Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. |

Images



Western Blotting

Image 1. Western Blot showing FBXO32 antibody used at a concentration of 1-2 ug/ml to detect its target protein.