

Datasheet for ABIN928987 **anti-IL17D antibody (N-Term)**



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	IL17D
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL17D antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	IL17 D antibody was raised in rabbit using the N terminal of IL17 as the immunogen
Purification:	Purified

Target Details

Target:	IL17D
Alternative Name:	IL17D (IL17D Products)
Background:	<p>The protein encoded by this gene is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines including IL6, IL8 and CSF2/ GM-CSF. The increased expression of IL8 induced by this cytokine was found to be NF-kappa B-dependent. Synonyms: Polyclonal IL17D antibody, Anti-IL17D antibody, interleukin 17D antibody, FLJ30846 antibody, IL-17D</p>

Target Details

antibody, IL-22 antibody, IL-27 antibody, IL27 antibody.

Pathways: [Cellular Response to Molecule of Bacterial Origin](#)

Application Details

Application Notes: WB: 0.2-1 $\mu\text{g}/\text{mL}$
Optimal conditions should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Concentration: Lot specific


Buffer: Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 $^{\circ}\text{C}$ /-20 $^{\circ}\text{C}$

Storage Comment: Store at 4 $^{\circ}\text{C}$, following reconstitution, aliquot and store at -20 $^{\circ}\text{C}$.

Images



Western Blotting

Image 1. IL17D antibody used at 0.2-1 $\mu\text{g}/\text{ml}$ to detect target protein.