

Datasheet for ABIN929186

anti-UCHL5 antibody (C-Term)





Go to Product page

| () | ve | V / | - | 1 A |
|--------|--------|------|---|-----|
| | \cup | 1 \/ | - | 1/1 |
| \sim | ' V C | 1 V | ı | v v |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | UCHL5 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This UCHL5 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | UCHL5 antibody was raised in rabbit using the C terminal of UCHL5 as the immunogen |
| Purification: | Purified |
| Target Details | |
| Target: | UCHL5 |
| Alternative Name: | UCHL5 (UCHL5 Products) |
| Background: | UCHL5 is the deubiquitinating enzyme associated with the proteasome. Synonyms: Polyclonal |
| | UCHL5 antibody, Anti-UCHL5 antibody, ubiquitin carboxyl-terminal hydrolase L5 antibody, CGI- |
| | 70 antibody, INO80R antibody, UCH37 antibody. |


Application Details

| Application Notes: | WB: 0.2-1 μg/mL |
|--------------------|--|
| | Optimal conditions should be determined by the investigator. |
| Comment: | UCHL5 Blocking Peptide, catalog no. 33R-4642, is also available for use as a blocking control in assays to test for specificity of this UCHL5 antibody |
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized | |
|------------------|---|--|
| Concentration: | Lot specific | |
| Buffer: | Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. | |

Images



Western Blotting

Image 1. Western Blot showing UCHL5 antibody used at a concentration of 1-2 ug/ml to detect its target protein.