

Datasheet for ABIN929380
anti-MAGEC2 antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

| | |
|----------------------|---------------------------------------|
| Quantity: | 100 µL |
| Target: | MAGEC2 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MAGEC2 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|---------------|--|
| Immunogen: | MAGEC2 antibody was raised in rabbit using the N terminal of MAGEC2 as the immunogen |
| Purification: | Purified |

Target Details

| | |
|-------------------|--|
| Target: | MAGEC2 |
| Alternative Name: | MAGEC2 (MAGEC2 Products) |
| Background: | This gene is a member of the MAGEC gene family. It is not expressed in normal tissues, except for testis, and is expressed in tumors of various histological types. This gene and the other MAGEC genes are clustered on chromosome Xq26-q27. Synonyms: Polyclonal MAGEC2 antibody, Anti-MAGEC2 antibody, melanoma antigen family C, 2 antibody, CT10 antibody, HCA587 antibody, MAGEE1 antibody, MGC13377 antibody. |

Application Details

| | |
|--------------------|--|
| Application Notes: | WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator. |
| Comment: | MAGEC2 Blocking Peptide, catalog no. 33R-8333, is also available for use as a blocking control in assays to test for specificity of this MAGEC2 antibody |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|--|
| Format: | Lyophilized |
| Concentration: | Lot specific |
| Buffer: | Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. |

Images



Western Blotting

Image 1. MAGEC2 antibody used at 0.2-1 ug/ml to detect target protein.