

# Datasheet for ABIN929450

# anti-CYP3A5 antibody (N-Term)





	e			

100 μL
CYP3A5
N-Term
Human
Rabbit
Polyclonal
This CYP3A5 antibody is un-conjugated
Western Blotting (WB)
CYP3 A5 antibody was raised in rabbit using the N terminal of CYP3 5 as the immunogen
Purified
CYP3A5
CYP3A5 (CYP3A5 Products)
This gene,CYP3A5, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some

#### **Target Details**

as well as the steroid hormones testosterone, progesterone and androstenedione. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. This cluster includes a pseudogene, CYP3A5P1, which is very similar to CYP3A5. This similarity has caused some difficulty in determining whether cloned sequences represent the gene or the pseudogene. Multiple alternatively spliced transcript variants have been identified for this gene. Synonyms: Polyclonal CYP3A5 antibody, Anti-CYP3A5 antibody, cytochrome P450, family 3, subfamily A, polypeptide 5 antibody, CP35 antibody, P450PCN3 antibody, PCN3 antibody.

Pathways:


Steroid Hormone Biosynthesis

### **Application Details**

Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	CYP3A5 Blocking Peptide, catalog no. 33R-1283, is also available for use as a blocking control in assays to test for specificity of this CYP3A5 antibody
Restrictions:	For Research Use only

### Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 $\mu$ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.



### **Western Blotting**

**Image 1.** Western Blot showing CYP3A5 antibody used at a concentration of 1-2 ug/ml to detect its target protein.