

Datasheet for ABIN934037  
**anti-FADD antibody (AA 1-208)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µL
Target:	FADD
Binding Specificity:	AA 1-208
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FADD antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

## Product Details

Immunogen:	FADD antibody was raised in mouse using recombinant human FADD (1-208aa) purified from E. coli as the immunogen.
Clone:	J1D2
Isotype:	IgG2b kappa
Purification:	FADD antibody was purified by protein-G affinity chromatography

## Target Details

Target:	FADD
Alternative Name:	FADD ( <a href="#">FADD Products</a> )
Pathways:	<a href="#">Apoptosis</a> , <a href="#">TLR Signaling</a> , <a href="#">Activation of Innate immune Response</a> , <a href="#">Positive Regulation of</a>

## Target Details

Endopeptidase Activity, Toll-Like Receptors Cascades

## Application Details

Application Notes: WB: 1:500-1:1000

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: Lot specific

Buffer: as a liquid PBS, pH 7.4, with 0.1 % NaN<sub>3</sub>.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid repeated freeze/thaw cycles

Storage: -20 °C

Storage Comment: Store at 4 °C for short term storage. Aliquot and store at -20 °C for long term storage.


## Images



**Human breast cancer tissue**

### Immunohistochemistry

**Image 1.** Paraffin embedded sections of human breast cancer tissue were incubated with nitrocellulose with anti-human FADD (1:50) for 2 hours at room temperature. Antigen retrieval was performed in 0.1M sodium citrate buffer and detected using Diaminobenzidine (DAB)



#### Western Blotting

**Image 2.** Cell lysates of Jurkat (30 ug) were resolved by SDS-PAGE, transferred to nitrocellulose membrane and probed with anti-human FADD (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.