

Datasheet for ABIN93932
anti-MME antibody (FITC)[Go to Product page](#)

4 Images

1 Publication

Overview

Quantity:	100 tests
Target:	MME
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MME antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	NALM-6 human pre-B cell line
Clone:	MEM-78
Isotype:	IgG1
Specificity:	The antibody MEM-78 reacts with an extracellular epitope CD10 antigen (CALLA - Common acute lymphatic leukemia antigen), a 100 kDa type II integral membrane protein.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	MME
---------	-----

Target Details

Alternative Name:	CD10 (MME Products)
Target Type:	Chemical
Background:	Membrane metalloendopeptidase,CD10 (neutral endopeptidase –, NEP, common acute lymphocytic leukemia antigen –, CALLA, membrane metallo-endopeptidase –, MME, enkephalinase) is a 100- kDa cell surface zinc metalloprotease, cleaving peptide bonds on the N-terminus of hydrophobic amino acids and inactivating multiple physiologically active peptides. CD10 is expressed on various normal cell types, including lymphoid precursor cells, germinal center B lymphocytes, and some epithelial cells, and its expression level serves as a marker for diagnostics of many carcinomas. CD10 is also a differentiation antigen for early B-lymphoid progenitors in the B-cell differentiation pathway and has a key role in regulation of growth, differentiation and signal transduction of many cellular systems.,CALLA, Neprilysin, Neutral endopeptidase, Enkephalinase, Atriopeptidase, MME
Gene ID:	4311
UniProt:	P08473
Pathways:	RTK Signaling , Peptide Hormone Metabolism , Regulation of Systemic Arterial Blood Pressure by Hormones , Smooth Muscle Cell Migration

Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Restrictions:	For Research Use only

Handling

Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

should be handled by trained staff only.

Handling Advice: **Do not freeze.**
Avoid prolonged exposure to light.


Storage: 4 °C

Storage Comment: Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Publications


Product cited in: Angelisová, Drbal, Horejsí, Cerný: "Association of CD10/neutral endopeptidase 24.11 with membrane microdomains rich in glycosylphosphatidylinositol-anchored proteins and Lyn kinase." in: **Blood**, Vol. 93, Issue 4, pp. 1437-9, (1999) ([PubMed](#)).

Images




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD10 (MEM-78) FITC antibody (20 µL reagent / 100 µL of peripheral whole blood).



Flow Cytometry

Image 2. Surface staining of NALM-6 human pre-B cell leukemia cell line with anti-human CD10 (MEM-78) PE. Total viable cells were used for analysis.



Flow Cytometry

Image 3. Surface staining of human peripheral blood with anti-human CD10 (MEM-78) FITC.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN93932.