

Datasheet for ABIN93946 anti-SEMA7A antibody (FITC)

1 Image

Publications

2



Overview

Quantity:	100 tests
Target:	SEMA7A
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SEMA7A antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	HPB-ALL human T cell line
Clone:	MEM-150
Isotype:	lgM
Specificity:	The antibody MEM-150 reacts with CD108 (JMH blood group antigen), a 80 kDa GPI-anchored extracellular glycoprotein expressed on various cell types including erythrocytes, lymphoblasts, at low levels it is present on circulating lymphocytes.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN93946 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	SEMA7A	
Alternative Name:	CD108 (SEMA7A Products)	
Background:	Semaphorin 7A (John Milton Hagen blood group),CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic cell migration, and in immune system affects cell proliferation, chemotaxis and cytokine release. On erythrocytes CD108 defines the JMH (John- Milton-Hagen) human blood group. CD108 signalizes through its receptors –, plexin C1 and beta1 integrins.,Semaphorin 7a, JMH blood group antigen, SEMAL, SEMAK1	
Gene ID:	8482	
UniProt:	075326	
Pathways:	Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Regulation of Cell Size	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.	
Comment:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.	

Handling

Restrictions:

Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
	Avoid prolonged exposure to light.


For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN93946 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

	•
land	lina
าสมณ	
ana	my

-	
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.
Publications	
Product cited in:	Angelisová, Drbal, Cerný, Hilgert, Horejsí: "Characterization of the human leukocyte GPI-
	anchored glycoprotein CDw108 and its relation to other similar molecules." in: Immunobiology,
	Vol. 200, Issue 2, pp. 234-45, (1999) (PubMed).
	Mudad, Rao, Angelisova, Horejsi, Telen: "Evidence that CDw108 membrane protein bears the
	JMH blood group antigen." in: Transfusion , Vol. 35, Issue 7, pp. 566-70, (1995) (PubMed).

Images



Flow Cytometry

Image 1. Surface staining of HPB-ALL human peripheral blood T cell leukemia cell line with anti-human CD108 (MEM-150) FITC. Total viable cells were used for analysis.