

Datasheet for ABIN93958
anti-CD11b antibody (Biotin)

4 Images

2 Publications

[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	CD11b (ITGAM)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD11b antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP)

Product Details

Immunogen:	Human granulocytes
Clone:	MEM-174
Isotype:	IgG2a
Specificity:	The antibody MEM-174 recognizes an extracellular epitope of CD11b antigen (Mac-1 alpha), a 165-170 kDa type I transmembrane protein mainly expressed on monocytes, granulocytes and NK-cells. The CD11b mediates neutrophil and monocyte interactions with stimulated endothelium.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and unconjugated antibody and free biotin are removed by size-exclusion chromatography.

Target Details

Target:	CD11b (ITGAM)
Alternative Name:	CD11b (ITGAM Products)
Background:	<p>Integrin subunit alpha M,CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.,Mac-1, Integrin alpha M, ITGAM, CR3A, MO1A, MAC1A</p>
Gene ID:	3684
UniProt:	P11215
Pathways:	Apoptosis , Activation of Innate immune Response , Toll-Like Receptors Cascades , Activated T Cell Proliferation

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-2 µg/mL, positive control: peripheral blood lymphocytes.
Comment:	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.
Storage:	4 °C

Handling


Storage Comment: Store at 2-8°C. Do not freeze.

Publications

Product cited in: Hasan, Osickova, Bumba, Novák, Sebo, Osicka: "Interaction of Bordetella adenylate cyclase toxin with complement receptor 3 involves multivalent glycan binding." in: **FEBS letters**, Vol. 589 , Issue 3, pp. 374-9, (2015) ([PubMed](#)).


Drbal, Moertelmaier, Holzhauser, Muhammad, Fuertbauer, Howorka, Hinterberger, Stockinger, Schütz: "Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement." in: **International immunology**, Vol. 19, Issue 5, pp. 675-84, (2007) ([PubMed](#)).

Images




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD11b (MEM-174) Biotin antibody (concentration in sample 6 µg/mL, Streptavidin APC).



Immunoprecipitation

Image 2. Immunoprecipitation of human CD11b/CD18 heterodimer from the lysate of washed PBMC isolated from healthy donor. Lysate was subjected to affinity column chromatography using anti-human CD11b (MEM-174) immunosorbent. Eluted immunoprecipitate was resolved on 7.5% SDS-PAGE and stained with Coomassie Blue.



Flow Cytometry

Image 3. Flow cytometry analysis (surface staining) of human peripheral blood with anti-human CD11b (MEM-174) biotin, streptavidin-APC.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN93958.