

Datasheet for ABIN93979  
**anti-CD147 antibody**



[Go to Product page](#)

2 Images

2 Publications

## Overview

Quantity:	0.1 mg
Target:	CD147 (BSG)
Reactivity:	Human, Pig
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD147 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	A soluble recombinant form of CD147, CD147Rg, which consists of the cDNA coding for the entire extracellular region of CD147 fused to the DNA coding for the hinge region, CH2 and CH3 domain of human IgG1.
Clone:	MEM-M6-2
Isotype:	IgG2b
Specificity:	The antibody MEM-M6/2 recognizes extracellular Ig domain D1 of CD147 (Neurothelin), a 50-60 kDa type I transmembrane glycoprotein primarily expressed on all leukocytes, red blood cells, platelets and endothelial cells, it is not expressed by resting lymphocytes. The antibody MEM-M6/1 is a high-affinity antibody capable of binding to unstimulated peripheral blood T cells.
Cross-Reactivity (Details):	Human, Porcine
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

## Target Details

Target:	CD147 (BSG)
Alternative Name:	CD147 ( <a href="#">BSG Products</a> )
Background:	Basigin (Ok blood group),CD147 (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumour cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays roles in cell development and activation, fetal development or retinal function.,Neurothelin, Basigin, LAM6, CSF, 5F7, TCSF, BSG, TCSF, Emprin, OK blood group
Gene ID:	682
UniProt:	<a href="#">P35613</a>
Pathways:	<a href="#">S100 Proteins</a>

## Application Details

Application Notes:	Flow cytometry: Recommended dilution: 4 µg/mL. Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/mL, positive tissue: testis.
Restrictions:	For Research Use only


## Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	<b>Do not freeze.</b>
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Product cited in: Plánka, Necas, Srnec, Rauser, Starý, Jancár, Amler, Filová, Hlucilová, Kren, Gál: "Use of allogenic stem cells for the prevention of bone bridge formation in miniature pigs." in: **Physiological research / Academia Scientiarum Bohemoslovaca**, Vol. 58, Issue 6, pp. 885-93, (2010) ([PubMed](#)).

Koch, Staffler, Hüttinger, Hilgert, Prager, Cerný, Steinlein, Majdic, Horejsí, Stockinger: "T cell activation-associated epitopes of CD147 in regulation of the T cell response, and their definition by antibody affinity and antigen density." in: **International immunology**, Vol. 11, Issue 5, pp. 777-86, (1999) ([PubMed](#)).

Images



Flow Cytometry

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD147 (MEM-M6/2) purified antibody (concentration in sample 0.6 µg/mL, GAM APC).

Flow Cytometry

**Image 2.** Separation of leukocytes stained using anti-human CD147 (MEM-M6/2) purified antibody (concentration in sample 0.6 µg/mL, GAM APC, red-filled) from leukocytes unstained by primary antibody (GAM APC, black-dashed) in flow cytometry analysis (surface staining).