
antibodies - online.com







anti-CD15 antibody



Images



\sim	
()\/\	rview
\cup	1 410 44

Quantity:	0.1 mg
Target:	CD15 (FUT4)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD15 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Cytometry by Time of Flight (CyTOF)

Product Details

Product Details	
Immunogen:	Human granulocytes
Clone:	MEM-158
Isotype:	IgM
Specificity:	The antibody MEM-158 reacts with CD15, a cell membrane molecule 3-fucosyl-N-acetylla ctosamine (3-FAL) strongly expressed on the surface of granulocytes, monocytes, macrophages, mast cells, it is also present on Langerhans cells and some myeloid precursors cells.
No Cross-Reactivity:	Cow, Pig, Sheep
Cross-Reactivity (Details):	Human
Purification:	Purified by sequential steps of physicochemical fractionation (differential precipitation and solid-phase chromatography methods).

Product Details

Purity:

> 95 % (by SDS-PAGE)

Target Details

Target: CD15 (FUT4)

Alternative Name: CD15 (FUT4 Products)

Background:

CD15 (Lewis x), also known as stage specific embryonic antigen-1 (SSEA-1) is a trisacharide determinant (3-fucosyl-N-acetyllactosamine) expressed on several glycolipids, glycoproteins and proteoglycans of various cell types, e.g. granulocytes, mast cells, monocytes, macrophages, cells of gastric mucosa, nervous system or various tumour cells. There are several structural relatives of Lewis x, e.g. sialyl-Lewis x or sulphated Lewis x. Cells with high surface expression of Le(x) antigen exhibit strong self-aggregation, based on calciumdependent Le(x)-Le(x) interaction. This process is involved for example in embryo compaction or in autoaggregation of teratocarcinoma cells. Sialyl-Le(x) and its isomer sialyl-Le(a) are ligands of selectins. CD15 expression has been extensively used to confirm diagnosis of Hodgkin's disease.,Lewis x Blood Group antigen, Le(x), SSEA-1, 3-fucosyl-N-acetyllactosamine

Application Details

Application Notes: Flow cytometry: Recommended dilution: 1-4 µg/mL.

Restrictions: For Research Use only

Handling

Concentration: 1 mg/mL

Buffer: Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Do not freeze.


Storage: 4 °C

Storage Comment: Store at 2-8°C. Do not freeze.




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD15 (MEM-158) purified antibody (concentration in sample 0,3 μ g/mL) GAM APC.



Flow Cytometry

Image 2. Flow Cytometry analysis Surface staining (flow cytometry) of human peripheral blood cells with anti-human CD15 (MEM-158) FITC. Cells in the granulocyte gate were used for analysis.



Flow Cytometry

Image 3. Surface staining (mass cytometry) of human peripheral blood cells (after ammonium chloride red blood cell lysis) with anti-human CD15 173Yb. Gated on singlets.

Please check the product details page for more images. Overall 5 images are available for ABIN93984.