antibodies - online.com







anti-CD16 antibody

Publications Images



<i>ا</i> ۱	1	-	M	/1	0	
u	1//	\vdash	1 \	/ I	-	\/\/
\sim	v	\sim	r٧		\sim	v v

Quantity:	0.1 mg	
Target:	CD16	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD16 antibody is un-conjugated	
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Western Blotting (WB)	

Product Details

Immunogen:	Human granulocytes
Clone:	MEM-154
Isotype:	lgG1
Specificity:	The antibody MEM-154 reacts with an extracellular epitope on CD16 antigen that is residing in proximity to FG loop (probably BC or C'E loop). CD16 is a low affinity receptor for aggregated IgG (FcgammaRIII antigen). The antibody MEM-154 reacts with CD16+ granulocytes, and it can be used for mapping CD16-158V/F polymorphism on NK cells, as it requires presence of V at amino acid 158.
Cross-Reactivity (Details):	Human
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	CD16
Alternative Name:	CD16 (CD16 Products)
Background:	CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms –, FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.,FcgammaRIII, IGFR3, FCRIII
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process
Application Details	
Application Notes:	Flow cytometry: Recommended dilution: 5-10 µg/mL, positive control: PBL (peripheral blood lymphocytes). The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. Western blotting: Non-reducing conditions.
Restrictions:	For Research Use only
Handling	
Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.
Publications	
Product cited in:	Gasdaska, Sherwood, Regan, Dickey: "An afucosylated anti-CD20 monoclonal antibody with


greater antibody-dependent cellular cytotoxicity and B-cell depletion and lower complement-dependent cytotoxicity than rituximab." in: **Molecular immunology**, Vol. 50, Issue 3, pp. 134-41, (2012) (PubMed).

Koene, Kleijer, Algra, Roos, von dem Borne, de Haas: "Fc gammaRIIIa-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gammaRIIIa, independently of the Fc gammaRIIIa-48L/R/H phenotype." in: **Blood**, Vol. 90, Issue 3, pp. 1109-14, (1997) (PubMed).

de Haas, Koene, Kleijer, de Vries, Simsek, van Tol, Roos, von dem Borne: "A triallelic Fc gamma receptor type IIIA polymorphism influences the binding of human IgG by NK cell Fc gamma RIIIa." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 156, Issue 8, pp. 2948-55, (1996) (PubMed).


Tamm, Schmidt: "The binding epitopes of human CD16 (Fc gamma RIII) monoclonal antibodies. Implications for ligand binding." in: **Journal of immunology (Baltimore, Md.: 1950)**, Vol. 157, Issue 4, pp. 1576-81, (1996) (PubMed).


Images



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD16 (MEM-154) purified antibody (concentration in sample $2\,\mu\text{g/mL}$) GAM APC.





Flow Cytometry

Image 2. Surface staining of human peripheral blood cells with anti-human CD16 (MEM-154) PE. The antibody MEM-154 does not react with CD16a present on NK cells in many subjects.

Flow Cytometry

Image 3. Surface staining of human peripheral blood cells with anti-human CD16 (MEM-154) purified, GAM/APC.