antibodies -online.com







Images



Go to Product page

\sim					
	1//	⊃r	V/I	Θ	Λ


Quantity:	100 tests	
Target:	CD177	
Reactivity:	Human, Non-Human Primate	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD177 antibody is conjugated to FITC	
Application:	Flow Cytometry (FACS)	

Product Details

Immunogen:	Human granulocytes
Clone:	MEM-166
Isotype:	lgG1
Specificity:	The antibody MEM-166 reacts with CD177 (Neutrophil specific antigen 1), a 60 kDa GPI-linked cell surface glycoprotein of uPAR family, expressed on granulocytes and in bone marrow early erythroblasts, megakaryocytes, promyelocytes and myelocytes.
Cross-Reactivity (Details):	Non-Human Primates, Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.


Target Details

rarget Details		
Target:	CD177	
Alternative Name:	CD177 (CD177 Products)	
Background:	CD177 Molecule,CD177 (NB1/HNA-2a and PRV-1 form) is a GPI-anchored glycoprotein present mainly on neutrophils. Its plasma membrane expression is increased during pregnancy and and inflammation or after G-CSF application. Ligand of CD177 has been identified as CD31 (PECAM-1). CD177 participates in neutrophil transmigration and seems to be also a proproliferative molecule. The antibodies against CD177 can be involved in neonatal alloimmune neutropenia (NAN).,NB1, PRV1, HNA2A	
Gene ID:	57126	
UniProt:	Q8N6Q3	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.	
Comment:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.	
Restrictions:	For Research Use only	
Handling		
Reconstitution:	No reconstitution is necessary.	
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD177 (MEM-166) FITC antibody (20 μ L reagent / 100 μ L of peripheral whole blood).



Flow Cytometry

Image 2. Surface staining of human peripheral blood cells with anti-CD177 (MEM-166) FITC.



Flow Cytometry

Image 3. Separation of human CD177 positive neutrophil granulocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD177 (MEM-166) FITC antibody (20 μ L reagent / 100 μ L of peripheral whole blood).