

Datasheet for ABIN94007

anti-Integrin beta 2 antibody (Biotin)

5 Images

5 Publications



[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	Integrin beta 2 (ITGB2)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Integrin beta 2 antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Leukocytes of a patient suffering from a LGL-type leukemia.
Clone:	MEM-48
Isotype:	IgG1
Specificity:	The antibody MEM-48 recognizes an extracellular epitope involving residues 534-546 in cysteine-rich repeat 3 of the CD18 antigen (integrin beta2 subunit, beta2 integrin). CD18 is a 90-95 kDa type I transmembrane protein expressed on all leukocytes.
No Cross-Reactivity:	Dog
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and unconjugated antibody and free biotin are removed by size-exclusion chromatography.

Target Details

Target:	Integrin beta 2 (ITGB2)
Alternative Name:	CD18 (ITGB2 Products)
Background:	<p>Integrin subunit beta 2,CD18, integrin beta2 subunit, forms heterodimers with four types of CD11 Molecule to constitute leukocyte (beta2) integrins: alphaLbeta2 (CD11a/CD18, LFA-1), alphaMbeta2 (CD11b/CD18, Mac-1, CR3), alphaXbeta2 (CD11c/CD18) and alphaDbeta2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts. Absence of CD18 leads to leukocyte adhesion deficiency-1, severe reduction of CD18 expression leads to the development of a psoriasiform skin disease. CD18 is also a target of Mannheimia (Pasteurella) haemolytica leukotoxin and is sufficient to mediate leukotoxin-mediated cytolysis.,Integrin beta2, Complement receptor C3 subunit beta, ITGB2</p>
Gene ID:	3689
UniProt:	P05107
Pathways:	NF-kappaB Signaling , Activation of Innate immune Response , Toll-Like Receptors Cascades , Activated T Cell Proliferation

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-3 µg/mL.
Comment:	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.




Handling

Storage: 4 °C

Storage Comment: Store at 2-8°C. Do not freeze.

Publications

- Product cited in: Kuttruff, Koch, Kelp, Pawelec, Rammensee, Steinle: "NKp80 defines and stimulates a reactive subset of CD8 T cells." in: **Blood**, Vol. 113, Issue 2, pp. 358-69, (2009) ([PubMed](#)).
- Garnotel, Rittié, Poitevin, Monboisse, Nguyen, Potron, Maquart, Randoux, Gillery: "Human blood monocytes interact with type I collagen through alpha x beta 2 integrin (CD11c-CD18, gp150-95)." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 164, Issue 11, pp. 5928-34, (2000) ([PubMed](#)).
- Stefanová, Horejsí: "Association of the CD59 and CD55 cell surface glycoproteins with other membrane molecules." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 147, Issue 5, pp. 1587-92, (1991) ([PubMed](#)).
- Bazil, Stefanová, Hilgert, Kristofová, Van?k, Horejsí: "Monoclonal antibodies against human leucocyte antigens. IV. Antibodies against subunits of the LFA-1 (CD11a/CD18) leucocyte-adhesion glycoprotein." in: **Folia biologica**, Vol. 36, Issue 1, pp. 41-50, (1990) ([PubMed](#)).
- Larson, Springer: "Structure and function of leukocyte integrins." in: **Immunological reviews**, Vol. 114, pp. 181-217, (1990) ([PubMed](#)).



Flow Cytometry

Image 1. Flow cytometry analysis (surface staining) of human peripheral blood with anti-CD18 (MEM-48) biotin, streptavidin-APC.

Western Blotting

Image 2. Fig. A Flow Cytometry analysis. Surface staining of human peripheral blood cells with anti-human CD18 (MEM-48) PE. Cells in the granulocyte gate were used for analysis. Fig. B Western Blotting

Immunohistochemistry

Image 3. Immunohistochemistry staining of human spleen (paraffin sections) using anti-CD18 (Commercially tested by LifeSpan BioSciences).

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN94007.