antibodies -online.com







anti-CD3 antibody (APC)

Images



Publications



Overview

Quantity:	100 tests
Target:	CD3
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD3 antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Human thymocytes and T lymphocytes.
Clone:	MEM-57
Isotype:	IgG2a kappa
Specificity:	The antibody MEM-57 reacts with an extracellular epitope on gamma-epsilon and delta-epsilon dimers of human CD3 complex, a part of a bigger multisubunit T cell receptor complex (CD3/TCR) expressed on peripheral blood T lymphocytes and mature thymocytes.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	CD3
Alternative Name:	CD3 (CD3 Products)
Background:	CD3 antigen, epsilon polypeptide,CD3 complex is crucial in transducing antigen-recognition
	signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR
	complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of
	the CD3 subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are
	structurally related members of the immunoglobulins super family encoded by closely linked
	genes on human chromosome 11. The CD3 components have long cytoplasmic tails that
	associate with cytoplasmic signal transduction molecules. This association is mediated at least
	in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may
	play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is
	present on 68-82 % of normal peripheral blood lymphocytes, 65-85 % of thymocytes and
	Purkynje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages
	of T lymphocytes may be observed in some autoimmune diseases.,CD3E, T3E, TCRE
Gene ID:	916
UniProt:	P07766
Pathways:	TCR Signaling, Ubiquitin Proteasome Pathway
Application Details	
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent
	/ 100 μL of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for
	100 tests.
Comment:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum
	conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct
	use. No reconstitution is necessary.
Restrictions:	For Research Use only
Handling	
	No reconstitution is necessary.
Reconstitution:	
Reconstitution: Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.
Publications	
Product cited in:	Blank, Welker, Haarer, Sterk, Nadalin, Yañez, Joos, Menrad, Snell, LaCorcia, Königsrainer,

roauct citea in:

Handgretinger, Schilbach: "Selective, efficient modulation of activated CD4+ ??T cells by the novel humanized antibody GZ-??TCR targeting human ??TCR." in: Bone marrow transplantation , (2014) (PubMed).

Majer, Vlaskova, Krol, Kalina, Kubanek, Stolnaya, Dvorakova, Elleder, Sikora: "Danon disease: a focus on processing of the novel LAMP2 mutation and comments on the beneficial use of peripheral white blood cells in the diagnosis of LAMP2 deficiency." in: Gene, Vol. 498, Issue 2, pp. 183-95, (2012) (PubMed).


Drbal, Moertelmaier, Holzhauser, Muhammad, Fuertbauer, Howorka, Hinterberger, Stockinger, Schütz: "Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement." in: International immunology, Vol. 19, Issue 5, pp. 675-84, (2007) (PubMed).

Brdicková, Brdicka, Angelisová, Horváth, Spicka, Hilgert, Paces, Simeoni, Kliche, Merten, Schraven, Horejsí: "LIME: a new membrane Raft-associated adaptor protein involved in CD4 and CD8 coreceptor signaling." in: The Journal of experimental medicine, Vol. 198, Issue 10, pp. 1453-62, (2003) (PubMed).

Panyi, Bagdány, Bodnár, Vámosi, Szentesi, Jenei, Mátyus, Varga, Waldmann, Gáspar, Damjanovich: "Colocalization and nonrandom distribution of Kv1.3 potassium channels and CD3 molecules in the plasma membrane of human T lymphocytes." in: Proceedings of the National Academy of Sciences of the United States of America, Vol. 100, Issue 5, pp. 2592-7, (2003) (PubMed).

There are more publications referencing this product on: Product page

Images




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD3 (MEM-57) APC antibody (10 μ L reagent / 100 μ L of peripheral whole blood).

Flow Cytometry

Image 2. Surface staining of human peripheral blood cells with anti-human CD3 (MEM-57) FITC. Cells in the lymphocyte gate were used for analysis.



CD3 APC

CD3 FITC

10⁴

105

0 10²

SS-A :: SS INT LIN

400K

200K

Flow Cytometry

Image 3. Flow cytometry analysis (surface staining) of CD3 in human peripheral blood with anti-CD3 (MEM-57) APC.

Please check the product details page for more images. Overall 4 images are available for ABIN94063.