

Datasheet for ABIN94125
anti-CD44 antibody (FITC)



[Go to Product page](#)

2 Images

3 Publications

Overview

Quantity:	100 tests
Target:	CD44
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD44 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Purpose:	Anti-Hu CD44 FITC
Immunogen:	Leukocytes of a patient suffering from LGL Type Leukaemia.
Clone:	MEM-85
Isotype:	IgG2b
Specificity:	The antibody MEM-85 reacts with an extracellular antigen of both cell surface-expressed and soluble form of CD44 antigen (Phagocyte glycoprotein 1), a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells and tissues (leukocytes, endothelial cells, mesenchymal cells, etc.), it is negative on platelets and hepatocytes.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	CD44
Alternative Name:	CD44 (CD44 Products)
Background:	CD44 Molecule,CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumour growth and metastasis, antiapoptosis and directional motility of cancer cells.,PGP-I, HUTCH-I, ECMR-III, Hermes antigen, Hyaluronate receptor, Heparan sulfate proteoglycan, Epican, MC56, MIC4, INLU, LHR
Gene ID:	960
UniProt:	P16070
Pathways:	Glycosaminoglycan Metabolic Process , Autophagy , Negative Regulation of intrinsic apoptotic Signaling

Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Restrictions:	For Research Use only

Handling

Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze. Avoid prolonged exposure to light.
Storage:	4 °C

Handling

Storage Comment: Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.


Publications

Product cited in: Kolar, Mehta, Pelayo, Capra: "A novel human B cell subpopulation representing the initial germinal center population to express AID." in: **Blood**, Vol. 109, Issue 6, pp. 2545-52, (2007) ([PubMed](#)).

Bazil, Strominger: "Metalloprotease and serine protease are involved in cleavage of CD43, CD44, and CD16 from stimulated human granulocytes. Induction of cleavage of L-selectin via CD16." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 152, Issue 3, pp. 1314-22, (1994) ([PubMed](#)).


Stefanová, Hilgert, Bazil, Kristofová, Horejsí: "Human leucocyte surface glycoprotein CDw44 and lymphocyte homing receptor are identical molecules." in: **Immunogenetics**, Vol. 29, Issue 6, pp. 402-4, (1989) ([PubMed](#)).

Images



Flow Cytometry

Image 1. Surface staining of human peripheral blood cells with anti-human CD44 (MEM-85) PE.



Flow Cytometry

Image 2. Surface staining of human peripheral blood with anti-human CD44 (MEM-85) FITC.