
# antibodies -online.com





# anti-CD55 antibody

**Images** 



Publication



## Overview

Quantity:	0.1 mg
Target:	CD55
Reactivity:	Human, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD55 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## **Product Details**


Immunogen:	HPB-ALL human T cell line
Clone:	MEM-118
Isotype:	IgM
Specificity:	The antibody MEM-118 recognizes an epitope in SCR4 domain of CD55 (Decay accelerating factor, DAF), a 60-70 kDa glycosylphosphatidylinositol (GPI)-anchored single chain extracellular glycoprotein. CD55 is widely expressed on hematopoietic and on many non-hematopoietic cells, it is weakly present on NK cells.
Cross-Reactivity (Details):	Human, Non-Human Primates
Purification:	Purified by sequential steps of physicochemical fractionation (differential precipitation and solid-phase chromatography methods).
Purity:	> 95 % (by SDS-PAGE)

# **Target Details**

Target:	CD55
Alternative Name:	CD55 (CD55 Products)
Background:	CD55 Molecule (Cromer blood group),CD55 (decay-accelerating factor, DAF) is a GPI-anchored
	membrane glycoprotein that protects autologous cells from classical and alternative pathway
	of complement cascade. Bidirectional interactions between CD55 and CD97 are involved in $T$
	cell regulation and CD55 can still regulate complement when bound to CD97. In tumours,
	besides protection agains complement, CD55 promotes neoangiogenesis, tumorigenesis,
	invasiveness and evasion of apoptosis.,DAF, CROM, CHAPLE
Gene ID:	1604
UniProt:	P08174
Pathways:	Complement System, Regulation of Leukocyte Mediated Immunity
Application Details	
Application Notes:	Flow cytometry: Recommended dilution: 10 μg/mL.
	Immunohistochemistry (paraffin sections): Recommended dilution: 10 μg/mL.
Restrictions:	For Research Use only
Handling	
Concentration:	1 mg/mL
Buffer:	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.
Publications	
Product cited in:	Angelisová, Drbal, Horejsí, Cerný: "Association of CD10/neutral endopeptidase 24.11 with
	membrane microdomains rich in glycosylphosphatidylinositol-anchored proteins and Lyn


kinase." in: **Blood**, Vol. 93, Issue 4, pp. 1437-9, (1999) (PubMed).

# **Images**



# **Flow Cytometry**

**Image 1.** Surface staining of human peripheral blood leukocytes by mouse monoclonal anti-CD55 antibody MEM-118.



# **Flow Cytometry**

**Image 2.** Surface staining of human peripheral blood cells with anti-CD55 (MEM-118) purified, GAM-APC.