

### Datasheet for ABIN94185

# anti-CD55 antibody (FITC)





Publication



Go to Product page

#### Overview

Quantity:	100 tests
Target:	CD55
Reactivity:	Human, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD55 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

#### **Product Details**

Purpose:	Anti-Hu CD55 FITC
Immunogen:	HPB-ALL human T cell line
Clone:	MEM-118
Isotype:	IgM
Specificity:	The antibody MEM-118 recognizes an epitope in SCR4 domain of CD55 (Decay accelerating factor, DAF), a 60-70 kDa glycosylphosphatidylinositol (GPI)-anchored single chain extracellular glycoprotein. CD55 is widely expressed on hematopoietic and on many non-hematopoietic cells, it is weakly present on NK cells.
Cross-Reactivity (Details):	Human, Non-Human Primates
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.


## **Target Details**

3	
Target:	CD55
Alternative Name:	CD55 (CD55 Products)
Background:	CD55 Molecule (Cromer blood group),CD55 (decay-accelerating factor, DAF) is a GPI-anchored membrane glycoprotein that protects autologous cells from classical and alternative pathway of complement cascade. Bidirectional interactions between CD55 and CD97 are involved in T cell regulation and CD55 can still regulate complement when bound to CD97. In tumours, besides protection agains complement, CD55 promotes neoangiogenesis, tumorigenesis, invasiveness and evasion of apoptosis.,DAF, CROM, CHAPLE
Gene ID:	1604
UniProt:	P08174
Pathways:	Complement System, Regulation of Leukocyte Mediated Immunity
Application Details	
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 20 $\mu$ L reagent / 100 $\mu$ L of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Restrictions:	For Research Use only
Handling	
Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.  Avoid prolonged exposure to light.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Product cited in:

Angelisová, Drbal, Horejsí, Cerný: "Association of CD10/neutral endopeptidase 24.11 with membrane microdomains rich in glycosylphosphatidylinositol-anchored proteins and Lyn kinase." in: **Blood**, Vol. 93, Issue 4, pp. 1437-9, (1999) (PubMed).

#### **Images**



#### **Flow Cytometry**

**Image 1.** Surface staining of human peripheral blood leukocytes by mouse monoclonal anti-CD55 antibody MEM-118.