

Datasheet for ABIN94187

anti-CD56 antibody**2** Images**4** Publications[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	CD56 (NCAM1)
Reactivity:	Human, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD56 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Immunogen:	KG-1 human acute myelogenous leukemia cell line
Clone:	MEM-188
Isotype:	IgG2a
Specificity:	The antibody MEM-188 reacts with an extracellular epitope on a 180 kDa isoform of CD56 (NCAM) expressed in leukocytes. It has been suggested that the antibody MEM-188 could react with rhesus monkey lymphocytes. Reactivity with other NCAM isoforms has not been tested.
Cross-Reactivity (Details):	Human, Non-Human Primates
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	CD56 (NCAM1)
Alternative Name:	CD56 (NCAM1 Products)
Background:	Neural cell adhesion molecule 1,CD56 (NCAM, neural cell adhesion molecule) is a transmembrane glycoprotein of immunoglobulin family serving as adhesive molecule which is ubiquitously expressed in nervous system, usually as 120 kDa, 140 kDa or 180 kDa isoform, and it is also found on T cells and NK cells. Polysialic modification results in reduction of CD56-mediated cell adhesion and is involved in cell migration, axonal growth, pathfinding and synaptic plasticity. CD56 is a widely used neuroendocrine marker with a high sensitivity for neuroendocrine tumours and ovarian granulosa cell tumours.,NCAM1, MSK39
Gene ID:	4684
UniProt:	P13591

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 3 µg/mL. Immunohistochemistry (paraffin sections): Recommended dilution: 1:50, pretreatment: heat retrieval (microwave oven) in citrate buffer, standard ABC visualization, positive tissue: membrane-specific reaction in adjacent peripheral nerve, staining of paraprostatic autonomic ganglion.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Publications

Product cited in:	Hovden, Karlsen, Jonsson, Aarstad, Appel: "Maturation of monocyte derived dendritic cells with
-------------------	--


OK432 boosts IL-12p70 secretion and conveys strong T-cell responses." in: **BMC immunology**, Vol. 12, pp. 2, (2011) ([PubMed](#)).

Olsen, Tollefsen, Aagaard, Reitan, Bannantine, Andersen, Sollid, Lundin: "Isolation of Mycobacterium avium subspecies paratuberculosis reactive CD4 T cells from intestinal biopsies of Crohn's disease patients." in: **PLoS ONE**, Vol. 4, Issue 5, pp. e5641, (2009) ([PubMed](#)).

Drbal, Moertelmaier, Holzhauser, Muhammad, Fuertbauer, Howorka, Hinterberger, Stockinger, Schütz: "Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement." in: **International immunology**, Vol. 19, Issue 5, pp. 675-84, (2007) ([PubMed](#)).


Yates, Rovis, Mitchell, Afzali, Tsang, Garin, Lechler, Lombardi, Garden: "The maintenance of human CD4+ CD25+ regulatory T cell function: IL-2, IL-4, IL-7 and IL-15 preserve optimal suppressive potency in vitro." in: **International immunology**, Vol. 19, Issue 6, pp. 785-99, (2007) ([PubMed](#)).

Images



Flow Cytometry

Image 1. Flow cytometry analysis (surface staining) of human peripheral blood lymphocytes with anti-CD56 (MEM-188) purified, GAM-APC.



Flow Cytometry

Image 2. Surface staining of human peripheral blood cells with anti-CD56 (MEM-188) FITC.