

Datasheet for ABIN94279
anti-Cytokeratin 18 antibody

8 Images

4 Publications

[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	Cytokeratin 18 (KRT18)
Reactivity:	Mammalian
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Cytokeratin 18 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Immunogen:	Cytoskeleton preparation of epidermal carcinoma cell line A431.
Clone:	C-04
Isotype:	IgG1
Specificity:	The antibody C-04 reacts with cytokeratin 18 (45 kDa intracellular antigen), a member of intermediate filaments subfamily represented in epithelial tissues.
Cross-Reactivity (Details):	Mammalian
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	Cytokeratin 18 (KRT18)
Alternative Name:	Cytokeratin 18 (KRT18 Products)
Background:	Keratin 18,Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families. Cytokeratin 18 belongs to type I family (acidic cytokeratins),K18, CK18, CYK18, KRT18
Gene ID:	3875
UniProt:	P05783
Pathways:	Apoptosis , Caspase Cascade in Apoptosis

Application Details

Application Notes:	Western blotting: Recommended dilution: 1-2 µg/mL. Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/mL, positive control: colon, kidney. ELISA: The antibody C-04 has been tested as the capture antibody in a sandwich ELISA for analysis of cytokeratin 18 in combination with the antibody DA-7 (cat. no. 1B-110-C100) and with the antibody DC-10 (cat. no. 1B-107-C100). Flow cytometry: Recommended dilution: 1-4 µg/mL. Intracellular staining.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Publications

Product cited in:


Bártek, Vojtšek, Stasková, Bártková, Kerekés, Rejthar, Kovarik: "A series of 14 new monoclonal antibodies to keratins: characterization and value in diagnostic histopathology." in: **The Journal of pathology**, Vol. 164, Issue 3, pp. 215-24, (1991) ([PubMed](#)).

Vojtšek, Stasková, Nenutil, Lauerová, Kovarik, Rejthar, Bártková, Bártek: "Monoclonal antibodies recognizing different epitopes of cytokeratin No.18." in: **Folia biologica**, Vol. 35, Issue 6, pp. 373-82, (1990) ([PubMed](#)).

Kovarik, Rejthar, Lauerová, Vojtšek, Bártková: "Monoclonal antibodies against individual cytokeratins in the detection of metastatic spread." in: **International journal of cancer. Supplement = Journal international du cancer. Supplement**, Vol. 3, pp. 50-5, (1989) ([PubMed](#)).


Lauerová, Kovarik, Bártek, Rejthar, Vojtšek: "Novel monoclonal antibodies defining epitope of human cytokeratin 18 molecule." in: **Hybridoma**, Vol. 7, Issue 5, pp. 495-504, (1989) ([PubMed](#)).

Images



Western Blotting

Image 1. Western blotting analysis of human cytokeratin 18 using mouse monoclonal antibody C-04 on lysates of HT-29 cell line and MOLT-4 cell line (cytokeratin non-expressing cell line, negative control) under non-reducing and reducing conditions. Nitrocellulose membrane was probed with 2 µg/mL of mouse anti-cytokeratin 18 monoclonal antibody followed by IRDye800-conjugated anti-mouse secondary antibody. A specific band was detected for cytokeratin 19 at approximately 46 kDa.



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded sections (kidney) Immunohistochemistry staining of human kidney (paraffin-embedded sections) with anti-Cytokeratin 18 (C-04).



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded sections (kidney) Immunohistochemistry staining of human kidney (paraffin-embedded sections) Fig. 3. Immunohistochemistry staining of human colon (paraffin-embedded sections) with anti-Cytokeratin 18 (Commercially tested by LifeSpan BioSciences).

Please check the [product details page](#) for more images. Overall 8 images are available for ABIN94279.