antibodies -online.com







anti-ATP6V1B1 antibody (C-Term)

Images



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Overview	
Quantity:	0.4 mL
Target:	ATP6V1B1
Binding Specificity:	AA 471-501, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATP6V1B1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	KLH conjugated synthetic peptide between 471-501 amino acids from the C-terminal region of human ATP6V1B1

Immunogen:	KLH conjugated synthetic peptide between 471-501 amino acids from the C-terminal region of human ATP6V1B1
Isotype:	lg Fraction
Specificity:	This antibody reacts to ATP6V1B1.
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Affinity chromatography on Protein A

Target Details

Target: ATP6V1B1


Target Details

Alternative Name:	ATP6V1B1 (ATP6V1B1 Products)	
Background:	This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that	
	mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle	
	acidification is necessary for such intracellular processes as protein sorting, zymogen	
	activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-	
	ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1	
	domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H	
	subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five	
	different subunits: a, c, c', c", and d. Additional isoforms of many of the V1 and V0 subunit	
	proteins are encoded by multiple genes or alternatively spliced transcript variants. This	
	encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney.	
	Mutations in this gene cause distal renal tubular acidosis associated with sensorineural	
	deafness. [provided by RefSeq].Synonyms: ATP6B1, Endomembrane proton pump 58 kDa	
	subunit, V-ATPase subunit B 1, V-type proton ATPase subunit B, VATB, VPP3, Vacuolar proton	
	pump subunit B 1, kidney isoform	
Gene ID:	525	
NCBI Accession:	NP_001683	
Pathways:	Sensory Perception of Sound, Transition Metal Ion Homeostasis, Proton Transport	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.25 mg/mL	
Buffer:	PBS containing 0.09 % (W/V) sodium azide as preservative	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	

Handling

Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. ATP6V1B1 Antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ATP6V1B1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Western Blotting

Image 2. ATP6V1B1 Antibody (C-term) western blot analysis in 293 cell line lysates (35μg/lane). This demonstrates the ATP6V1B1 antibody detected the ATP6V1B1 protein (arrow).