



[Go to Product page](#)

Datasheet for ABIN950972

anti-C11orf95 antibody (C-Term)

3 Images

Overview

Quantity:	0.4 mL
Target:	C11orf95
Binding Specificity:	AA 563-592, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C11orf95 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 563-592 amino acids from the C-terminal region of human C9JLR9
Isotype:	Ig Fraction
Specificity:	This antibody reacts to C9JLR9.
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Affinity chromatography on Protein A

Target Details

Target:	C11orf95
---------	----------

Target Details

Abstract:	C11orf95 Products
Background:	Synonyms: C11orf95, Uncharacterized protein C11orf95
Gene ID:	65998
NCBI Accession:	NP_001138408


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) sodium azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

Images



Immunofluorescence

Image 1. Confocal immunofluorescent analysis of C9JLR9 Antibody (C-term)(Cat#AP50677PU-N) with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. C9JLR9 Antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human adipose tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of C9JLR9 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 3. C9JLR9 Antibody (C-term) western blot analysis in HepG2 cell line lysates (35µg/lane). This demonstrates the C9JLR9 antibody detected the C9JLR9 protein (arrow).