antibodies - online.com







anti-CLEC3B antibody (Middle Region)

Images



\sim	
()\/\	rview
\cup	

Quantity:	0.4 mL
Target:	CLEC3B
Binding Specificity:	AA 102-130, Middle Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLEC3B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	KLH conjugated synthetic peptide between 102-130 amino acids from the Central region of human CLEC3B
Isotype:	lg Fraction
Specificity:	This antibody reacts to CLEC3B.
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Saturated Ammonium Sulfate (SAS) precipitation
Target Details	
Target:	CLEC3B

Target Details

Alternative Name:	CLEC3B (CLEC3B Products)
Background:	Tetranectin binds to plasminogen and to isolated kringle 4. It may be involved in the packaging of molecules destined for exocytosis. Synonyms: C-type lectin domain family 3 member B, Plasminogen kringle 4-binding protein, TNA, Tetranectin
Molecular Weight:	22567 Da
Gene ID:	7123
NCBI Accession:	NP_003269


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS, 0.09 % (W/V) sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.





Immunohistochemistry (Paraffin-embedded Sections)

Image 1. CLEC3B Antibody (Center) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CLEC3B Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Western Blotting

Image 2. Western blot analysis of CLEC3B (arrow) using rabbit polyclonal CLEC3B Antibody (Center) . 293 cell lysates (2 μ g/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CLEC3B gene.