

Datasheet for ABIN952260

anti-FAM96B antibody (C-Term, N-Term)**3** Images[Go to Product page](#)

Overview

| | |
|----------------------|---|
| Quantity: | 0.4 mL |
| Target: | FAM96B |
| Binding Specificity: | C-Term, N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FAM96B antibody is un-conjugated |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA) |

Product Details

| | |
|---------------|---|
| Immunogen: | KLH conjugated synthetic peptide from the C-terminal region of human FAM96B |
| Isotype: | Ig Fraction |
| Specificity: | This antibody recognizes Human FAM96B (N-term). |
| Purification: | Affinity Chromatography on Protein A |

Target Details

| | |
|-------------------|--|
| Target: | FAM96B |
| Alternative Name: | FAM96B (FAM96B Products) |
| Background: | Synonyms: CGI-128, HSPC118 |

Target Details

| | |
|-------------------|---------------------------|
| Molecular Weight: | 17663 Da |
| Gene ID: | 51647 |
| NCBI Accession: | NP_057146 |


Application Details

| | |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling


| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 0.25 mg/mL |
| Buffer: | PBS containing 0.09 % (W/V) Sodium Azide as preservative |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Avoid repeated freezing and thawing. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer. |

Images




Flow Cytometry

Image 1. Flow cytometric analysis of A375 cells using FAM96B Antibody (N-term) Cat.-No AP51505PU-N (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-fixed and paraffin-embedded human skin reacted with FAM96B Antibody (N-term) followed which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.



Western Blotting

Image 3. Western blot analysis of FAM96B Antibody (N-term) in A375 cell line lysates (35ug/lane). FA96B (arrow) was detected using the purified Pab.