

Datasheet for ABIN952571  
**anti-GOLGA8A antibody (N-Term)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	0.4 mL
Target:	GOLGA8A
Binding Specificity:	AA 136-164, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

## Product Details

Immunogen:	KLH conjugated synthetic peptide between 136-164 amino acids from the N-terminal region of human GOLGA8A.
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human GOLGA8A (N-term).
Purification:	Protein A column, followed by peptide affinity purification

## Target Details

Target:	GOLGA8A
Abstract:	<a href="#">GOLGA8A Products</a>
Background:	Synonyms: GOLGA8A, Golgin subfamily A member 8A

## Target Details

Molecular Weight:	51kd
Gene ID:	23015
NCBI Accession:	<a href="#">NP_851422</a>


## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

## Images




### Immunofluorescence

**Image 1.** Confocal immunofluorescent analysis of GOG8A Antibody (N-term) Cat.-No AP51887PU-N with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue reacted with GOG8A Antibody (N-term) followed by peroxidase conjugation of the secondary antibody and DAB staining.



#### Western Blotting

**Image 3.** Western blot analysis of GOG8A Antibody (N-term) in 293 cell line lysates (35ug/lane). This demonstrates the GOG8A antibody detected the GOG8A protein (arrow).