

Datasheet for ABIN952687  
**anti-HCFC1R1 antibody (N-Term)**

## 2 Images

[Go to Product page](#)

## Overview

Quantity:	0.4 mL
Target:	HCFC1R1
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HCFC1R1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

## Product Details

Immunogen:	KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human HCFC1R1.
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human HCFC1R1 / HPIP (N-term).
Purification:	Protein A column, followed by peptide affinity purification

## Target Details

Target:	HCFC1R1
Alternative Name:	HCFC1R1 / HPIP ( <a href="#">HCFC1R1 Products</a> )

## Target Details


Background:	Regulates HCFC1 activity by modulating its subcellular localization. Overexpression of HCFC1R1 leads to accumulation of HCFC1 in the cytoplasm. HCFC1R1-mediated export may provide the pool of cytoplasmic HCFC1 required for import of virion-derived VP16 into the nucleus.Synonyms: Host cell factor C1 regulator 1
Molecular Weight:	15291 Da
Gene ID:	54985
NCBI Accession:	<a href="#">NP_001002017</a>

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



**Immunohistochemistry (Paraffin-embedded Sections)**

**Image 1.** Immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue reacted with HCFC1R1 / HPIP Antibody (N-term) followed by peroxidase conjugation of the secondary antibody and DAB staining.



**Western Blotting**

**Image 2.** Western blot analysis of HCFC1R1 / HPIP Antibody (N-term) in MDA-MB231 cell line lysates (35ug/lane).