



Datasheet for ABIN953361
anti-MDM1 antibody (C-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	0.4 mL
Target:	MDM1
Binding Specificity:	AA 655-684, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 655-684 amino acids from the C-terminal region of human MDM1
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human MDM1 (C-term).
Purification:	Protein A column, followed by peptide affinity purification

Target Details

Target:	MDM1
Alternative Name:	MDM1 (MDM1 Products)
Background:	This gene encodes a nuclear protein similar to the mouse double minute 1 protein. The mouse gene is located in double minute (DM) chromatin particles and is amplified in the mouse

Target Details

transformed 3T3 cell line, and the protein is able to bind to p53. In mouse several transcripts have been described for this gene which result from alternative polyadenylation, splicing and exon usage. Synonyms: Nuclear protein MDM1

Molecular Weight: 80735 Da

Gene ID: 56890

NCBI Accession: [NP_001191957](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.25 mg/mL

Buffer: PBS containing 0.09 % (W/V) Sodium Azide as preservative


Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid repeated freezing and thawing.


Storage: 4 °C/-20 °C

Storage Comment: Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue reacted with MDM1 Antibody (C-term) followed which was peroxidase conjugated to the secondary antibody and followed by DAB staining. This data demonstrates the use of MDM1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of MDM1 Antibody (C-term) in A549 cell line lysates (35ug/lane). This demonstrates the MDM1 antibody detected the MDM1 protein (arrow).