

Datasheet for ABIN953548  
**anti-MMRN2 antibody (C-Term)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	0.4 mL
Target:	MMRN2
Binding Specificity:	AA 767-797, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MMRN2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

## Product Details

Immunogen:	KLH conjugated synthetic peptide between 767-797 amino acids from the C-terminal region of Human Multimerin-2 Genename: MMRN2
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human Multimerin-2 (C-term).
Purification:	Protein A column, followed by peptide affinity purification

## Target Details

Target:	MMRN2
Alternative Name:	Multimerin-2 ( <a href="#">MMRN2 Products</a> )

## Target Details


Background:	Synonyms: EMILIN-3, EMILIN3, Elastin microfibril interface located protein 3, Elastin microfibril interfacier 3, EndoGlyx-1 p125/p140 subunit, MMRN2
Molecular Weight:	104409 Da
Gene ID:	79812
NCBI Accession:	<a href="#">NP_079032</a>

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue reacted with Multimerin-2 Antibody (C-term) followed by peroxidase conjugation of the secondary antibody and DAB staining.



#### Western Blotting

**Image 2.** Western blot analysis of Multimerin-2 Antibody (C-term) in NCI-H460 cell line lysates (35ug/lane). This demonstrates the MMRN2 antibody detected the MMRN2 protein (arrow).