antibodies - online.com








anti-OR4F5 antibody (N-Term)







\sim	
()\/△	rview
\cup	1 410 44

Background:

Quantity:	0.4 mL
Target:	OR4F5
Binding Specificity:	AA 60-90, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR4F5 antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	KLH conjugated synthetic peptide between 60-90 amino acids from the N-terminal region of
	human OR4F5
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human OR4F5 (N-term).
Purification:	Protein A column, followed by peptide affinity purification
Target Details	
Target:	OR4F5
Alternative Name:	OR4F5 (OR4F5 Products)

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response

that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. Synonyms: Olfactory receptor 4F5

Molecular Weight: 34330 Da

Gene ID: 79501

NCBI Accession: NP_001005484


Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



Western Blotting

Image 1. Western blot analysis of OR4F5 Antibody (Nterm) in A549 cell line lysates (35ug/lane). This demonstrates the OR4F5 antibody detected the OR4F5 protein (arrow).