



[Go to Product page](#)

Datasheet for ABIN953926
anti-OR5L2 antibody (N-Term)

2 Images

Overview

Quantity:	0.4 mL
Target:	OR5L2
Binding Specificity:	AA 62-90, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR5L2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 62-90 amino acids from the N-terminal region of human OR5L2
Isotype:	Ig Fraction
Specificity:	This antibody recognizes Human OR5L2 (C-term).
Purification:	Protein A column, followed by peptide affinity purification

Target Details

Target:	OR5L2
Alternative Name:	OR5L2 (OR5L2 Products)

Target Details

Background: Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. Synonyms: Olfactory receptor 5L2

Molecular Weight: 34682 Da

Gene ID: 26338

NCBI Accession: [NP_001004739](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.25 mg/mL

Buffer: PBS containing 0.09 % (W/V) Sodium Azide as preservative


Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid repeated freezing and thawing.


Storage: 4 °C/-20 °C

Storage Comment: Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue reacted with OR5L2 Antibody (N-term) followed which was peroxidase conjugated to the secondary antibody and followed by DAB staining. This data demonstrates the use of the OR5L2 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of OR5L2 Antibody (N-term) in CEM, HeLa cell line lysates (35ug/lane). This demonstrates the OR5L2 antibody detected the OR5L2 protein (arrow).