# antibodies - online.com







### anti-PCDHB14 antibody (N-Term)

**Images** 




$\sim$	
( )\/\	rview
$\cup$	

Quantity:	0.4 mL
Target:	PCDHB14
Binding Specificity:	AA 165-193, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDHB14 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	Synthetic peptide - KLH conjugated - corresponding to the N-terminal region (between 165-193aa) of human PCDHB14.
Isotype:	Ig Fraction
Specificity:	This antibody detects human PCDHB14 at N-term.
Cross-Reactivity (Details):	Species reactivity (tested):Human
Purification:	Purified through a protein A column; followed by peptide affinity purification.
Target Details	
Target:	PCDHB14

### **Target Details**

PCDHB14 (PCDHB14 Products)	
The PCDHB14 gene is a member of the protocadherin beta gene cluster, one of three related	
gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusua	
genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluste	
contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a	
cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular	
domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the	
alpha and gamma clusters, the transcripts from these genes are made up of only one large	
exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion	
proteins are integral plasma membrane proteins. Their specific functions are unknown but they	
most likely play a critical role in the establishment and function of specific cell-cell neural	
connections.Synonyms: PCDH-beta-14, Protocadherin beta-14	
56122	
NP_061757	
Optimal working dilution should be determined by the investigator.	
For Research Use only	
Liquid	
0.25 mg/mL	
PBS with 0.09 % (W/V) Sodium Azide.	
Sodium azide	
This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
should be handled by trained staff only.	
Avoid repeated freezing and thawing.	
4 °C/-20 °C	
Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.	



## 250 130 95 - 4 72 - 55

### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemistry analysis in human brain tissue (formalin-fixed, paraffin-embedded) using PCDHB14 Antibody (N-term), followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PCDHB14 antibody for IHC. Clinical relevance has not been evaluated.

#### **Western Blotting**

**Image 2.** Western blot analysis in K562 cell line lysates (35ug/lane) using PCDHB14 Antibody (N-term). This demonstrates the PCDHB14 antibody detected the PCDHB14 protein (arrow).