antibodies -online.com





anti-PRPF38A antibody (C-Term)

3 Images



Go to Product page

\sim					
	1//6	r	V I	Θ	Λ

Quantity:	0.4 mL	
Target:	PRPF38A	
Binding Specificity:	AA 283-312, C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PRPF38A antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded	
	Sections) (IHC (p)), Enzyme Immunoassay (EIA)	
Product Details		
Immunogen:	KLH conjugated synthetic peptide between 283~312 amino acids from the C-terminal region of Human PRPF38A	
Isotype:	lg Fraction	
Specificity:	This antibody recognizes Human PRPF38A (C-term).	
Purification:	Affinity Chromatography on Protein A	
Target Details		
Target:	PRPF38A	
Alternative Name:	PRPF38A (PRPF38A Products)	

Target Details

Background:	Synonyms: Pre-mRNA-splicing factor 38A	
Molecular Weight:	37477 Da	
Gene ID:	84950	
NCBI Accession:	NP_116253	


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Format:	Liquid	
Concentration:	0.25 mg/mL	
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.	

Images




Flow Cytometry

Image 1. Flow cytometric analysis of HepG2 cells using PRPF38A Antibody (C-term) Cat.-No AP53428PU-N (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin fixed and paraffin embedded human cervix carcinoma reacted with PRPF38A Antibody (Cterm) followed by peroxidase conjugation of the secondary antibody and DAB staining



Western Blotting

Image 3. Western blot analysis of PRPF38A Antibody (Cterm) in MDA-MB231,CEM,HL-60 cell line lysates (35ug/lane). This demonstrates the PR38A antibody detected the PR38A protein (arrow).