

Datasheet for ABIN955354
anti-TSGA10IP antibody (C-Term)

2 Images

[Go to Product page](#)

Overview

Quantity:	0.4 mL
Target:	TSGA10IP
Binding Specificity:	AA 499-528, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TSGA10IP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 499-528 amino acids from the C-terminal region of human TSGA10IP
Isotype:	Ig Fraction
Specificity:	This antibody detects TSGA10IP (C-term).
Cross-Reactivity (Details):	Species reactivity (tested):Human
Purification:	Protein A column followed by peptide affinity purification

Target Details

Target:	TSGA10IP
---------	----------

Target Details

Alternative Name:	TSGA10IP (TSGA10IP Products)
Background:	Synonyms: Testis-specific protein 10-interacting protein, Tsga10-interacting protein
Gene ID:	254187
NCBI Accession:	NP_689975

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS with 0.09 % (W/V) sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2 - 8 °C for up to six months or (in aliquots) at -20 °C for longer.

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TSGA10IP Antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TSGA10IP Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. TSGA10IP Antibody (C-term) western blot analysis in Ramos cell line lysates (35 µg/lane). This demonstrates the TSGA10IP antibody detected the TSGA10IP protein (arrow).