



[Go to Product page](#)

Datasheet for ABIN961196
anti-PBX3 antibody (AA 342-435)

2 Images

Overview

| | |
|----------------------|---|
| Quantity: | 50 µg |
| Target: | PBX3 |
| Binding Specificity: | AA 342-435 |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This PBX3 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|---------------|---|
| Brand: | IHC-plus™ |
| Immunogen: | PBX3 (NP_006186, 342 a.a. ~ 435 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. Type of Immunogen: Recombinant protein |
| Clone: | 1A11 |
| Isotype: | IgG2a kappa |
| Specificity: | Human Pbx3 |
| Purification: | Purified from ascites by Protein A |

Target Details

Target: PBX3

Alternative Name: PBX3 ([PBX3 Products](#))

Background: Name/Gene ID: PBX3

Synonyms: PBX3, Pre-B-cell leukemia homeobox 3, Homeobox protein PBX3

Gene ID: 5090

NCBI Accession: [NP_006186](#)

UniProt: [P40426](#)

Application Details

Application Notes: Approved: ELISA, IHC, IHC-P (5 µg/mL), WB

Usage: Immunohistochemistry: Formalin-fixed paraffin-embedded sections. Sandwich ELISA: Recombinant protein. Western Blot: Cell lysate, Recombinant protein.

Comment: Target Species of Antibody: Human

Restrictions: For Research Use only

Handling

Format: Liquid


Concentration: Lot specific

Buffer: PBS, pH 7.4

Handling Advice: Aliquot to Avoid repeated freezing and thawing.


Storage: -20 °C

Storage Comment: Store at -20°C or lower. Aliquot to avoid freeze-thaw cycles.



Immunohistochemistry

Image 1. Anti-PBX3 antibody IHC staining of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN961196 concentration 5 ug/ml.



Immunohistochemistry

Image 2. Anti-PBX3 antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.