

Datasheet for ABIN964243

**Goat IgG isotype control (Rhodamine)****2** Images[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	IgG
Host:	Goat
Antibody Type:	Native
Conjugate:	Rhodamine
Application:	Isotype Control (IsoC), ELISA, Western Blotting (WB), Immunomicroscopy (IM)

## Product Details

Isotype:	IgG
Fragment:	Fc fragment
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

## Target Details

Target:	IgG
Abstract:	<a href="#">IgG Products</a>
Target Type:	Antibody

## Application Details

Application Notes:	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
--------------------	---

## Application Details

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Restore with deionized water (or equivalent)

Concentration: 1.0 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2


Preservative: Sodium azide

Precaution of Use: **WARNING:** Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Product is photosensitive and should be protected from light.


Storage: 4 °C

## Validation report #103142 for Immunofluorescence (IF)



### Dot Blot

**Image 1.** Dot Blot of Rhodamine Conjugated Goat IgG Fc. Dotted Rhodamine Conjugated Goat IgG Fc with following concentrations. Load: Lane 1 - 50ng Lane 2 - 16.67ng Lane 3 - 5.56ng Lane 4 - 1.85ng Lane 5 - 0.62ng Primary antibody: none Secondary antibody: none Block: ABIN925618 for 1 HR at RT.



SDS-PAGE

**Image 2.** SDS-PAGE of Goat IgG F(c) Fragment Rhodamine Conjugated . Lane M: 5 µL Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule . Lane 2: Reduced Goat IgG F(c) Fragment Rhodamine Conjugated . Lane 3: Reduced Goat IgG F(ab) Fragment . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 µg for IgG, F(c) and F(ab); 3 µg for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.