



Datasheet for ABIN964252

Goat IgG isotype control (FITC)



[Go to Product page](#)

1 Image

Overview

| | |
|----------------|---|
| Quantity: | 1 mg |
| Target: | IgG |
| Host: | Goat |
| Antibody Type: | Native |
| Conjugate: | FITC |
| Application: | Isotype Control (IsoC), ELISA, Western Blotting (WB), Immunomicroscopy (IM) |

Product Details

| | |
|------------------|--|
| Isotype: | IgG |
| Characteristics: | Concentration Definition: by UV absorbance at 280 nm |

Target Details

| | |
|--------------|------------------------------|
| Target: | IgG |
| Abstract: | IgG Products |
| Target Type: | Antibody |

Application Details

| | |
|--------------------|---|
| Application Notes: | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| Comment: | Excitation/Emission wavelength: 494 nm/514 nm |

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Restore with deionized water (or equivalent)

Concentration: 1.0 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2


Preservative: Sodium azide

Precaution of Use: **WARNING:** Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Product is photosensitive and should be protected from light.

Storage: 4 °C

Images



SDS-PAGE

Image 1. SDS-PAGE of Goat IgG Whole Molecule Fluorescein Conjugated . Lane M: 5 μ L Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule Fluorescein Conjugated . Lane 2: Reduced Goat IgG F(c) Fragment . Lane 3: Reduced Goat IgG F(ab) Fragment . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 μ g for IgG, F(c) and F(ab); 3 μ g for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.