

Datasheet for ABIN964831

Rabbit anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (FITC)



[Go to Product page](#)

Overview

Quantity:	20 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	FITC
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)

Product Details

Purpose:	F(ab') ₂ Guinea Pig IgG (H&L) Antibody Fluorescein Conjugated
Immunogen:	Optional[Immunogen]: Guinea Pig IgG whole molecule
Isotype:	IgG
Fragment:	F(ab') ₂ fragment
Cross-Reactivity (Details):	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.
Characteristics:	F(ab') ₂ Anti-Guinea Pig IgG F(ab') ₂ Antibody generated in rabbit detects Guinea Pig F(ab') ₂ . Representing approximately 75 % of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types.

Product Details

When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Purification: This product is a F(ab')₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above.

Target Details

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

Background: Because of their smaller size, F(ab)₂ fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)₂ fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)₂ fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Application Details

Application Notes: Application Note: This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Flow Cytometry Dilution: 1:500 - 1:2,500 FLISA Dilution: 1:10,000 - 1:50,000 IF Microscopy Dilution: 1:1,000 - 1:5,000 Other: FLOW CYTOMETRY 1:500 - 1:2,500

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Buffer: Restore with deionized water (or equivalent), Reconstitution Volume: 2.0 mL
Concentration:	10.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free , Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months