antibodies -online.com



Datasheet for ABIN965005

Goat anti-Rat IgG (Heavy & Light Chain) Antibody (Atto 594) - Preadsorbed



Go to Product page

2 Images

Overview	
Quantity:	100 μg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Atto 594
Application:	Western Blotting (WB), FLISA, Fluorescence Microscopy (FM)

Product Details


Immunogen: Rat IgG whole molecule
IgG
Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG and Rat Serum.
Anti-Rat IgG (H&L) conjugated to ATTO 594 is designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. This product is designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial
suitable for multiplex analysis, including multicolor imaging, utilizing various commercial

Product Details

1 Toddet Details	
	platforms.
Purification:	Preadsorption: Solid phase absorption
Labeling Ratio:	2.5
Target Details	
Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Goat anti-Rat IgG ATTO594 Conjugated Antibody, Goat anti-Rat IgG Antibody ATTO
	594 Conjugation
	Background: Anti-Rat IgG (H&L) conjugated to ATTO 594 is designed for STED microscopy,
	FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and
	fluorescent western blotting. This product is also suitable for multiplex analysis, including
	multicolor imaging, utilizing various commercial platforms.
Application Details	
Application Notes:	Application Note: The emission spectra for this ATTO conjugate matches the principle output
	wavelengths of most common fluorescence instrumentation.
	FLISA Dilution: >1:20,000
	Western Blot Dilution: >1:10,000
	IF Microscopy Dilution: >1:5,000
Comment:	The emission spectra for this ATTO conjugate matches the principle output wavelengths of
	most common fluorescence instrumentation.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 500 μL
	Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2


	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at 4 °C prior to restoration. For extended storage aliquot contents and freeze at -24 °C or below. This product is stable for several weeks at 4 °C as an undiluted liquid.
Expiry Date:	12 months

Validation report #103828 for ELISA (ELISA)



Western Blotting

Image 1. Western Blot of Anti-Rat IgG (H&L) (GOAT) Antibody (Min X Bv Ch Gt GP Ham Hs Hu Ms Rb & Sh Serum Proteins). Lane M: 3 µl Molecular Ladder. Lane 1: Rat IgG whole molecule. Lane 2: Rat IgG F(c) Fragment. Lane 3: Rat IgG Fab Fragment. Lane 4: Rat IgM Whole Molecule. Lane 5: Rat Serum. All samples were reduced. Load: 50 ng per Iane. Block: ABIN925618 for 30 min at RT. Primary Antibody: Anti-Rat IgG (H&L) (GOAT) Antibody (Min X Bv Ch Gt GP Ham Hs Hu Ms Rb & Sh Serum Proteins) 1:1,000 for 60 min at RT. Secondary Antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in ABIN925618 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and Fab, 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.



Immunofluorescence

Image 2.