

Datasheet for ABIN965059

## Goat anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (Biotin) - Preadsorbed



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

### Product Details

Immunogen:	Immunogen: Guinea Pig IgG whole molecule
Isotype:	IgG
Fragment:	F(ab') <sub>2</sub> fragment
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum.
Purification:	Preadsorption: Solid phase absorption
Sterility:	Sterile filtered

### Target Details

Target:	IgG
---------	-----

## Target Details

---

Abstract: [IgG Products](#)

---

Target Type: Antibody

---

Background: Synonyms: Goat F(ab')<sub>2</sub> Anti-Guinea Pig IgG Antibody Biotin Conjugation, Goat F(ab')<sub>2</sub> Anti-Guinea Pig IgG Biotin Conjugated Antibody, Goat Fab2 Anti-Guinea Pig IgG Biotin Conjugated Antibody

Background: F(ab')<sub>2</sub> Anti-Guinea Pig IgG Biotin Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)<sub>2</sub> fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)<sub>2</sub> fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)<sub>2</sub> fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')<sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

## Application Details

---

Application Notes: Immunohistochemistry Dilution: 1:1,000 - 1:5,000

Application Note: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 µg of Guinea Pig IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 is suggested for this product.

ELISA Dilution: 1:20,000 - 1:100,000

Western Blot Dilution: 1:2,000 - 1:10,000

---

Comment: Post Translational Modification: Phosphorylation

---

Restrictions: For Research Use only

---

## Handling

---

Format: Lyophilized

---

Reconstitution: Reconstitution Volume: 1.0 mL  
Reconstitution Buffer: Restore with deionized water (or equivalent)

---

Concentration: 1.0 mg/mL

---

## Handling

---

Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below.
Storage:	RT, 4 °C, -20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening. Store the vial at -20 °C or below after dilution.
Expiry Date:	12 months