

### Datasheet for ABIN965060

# Goat anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (Texas Red (TR)) - Preadsorbed

1 mg



Go to Product page

Overviev	

Quantity:

Quartity.	
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Texas Red (TR)
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)
Product Details	
Immunogen:	Immunogen: Guinea Pig IgG whole molecule
Isotype:	lgG
Isotype: Fragment:	IgG F(ab')2 fragment
Fragment:	F(ab')2 fragment  Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum,
Fragment: Specificity:	F(ab')2 fragment  Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum.  This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex
Fragment: Specificity: Characteristics:	F(ab')2 fragment  Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum.  This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

### Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Goat F(ab')2 Anti-Guinea Pig IgG Antibody Texas Red™ Conjugation, Goat F(ab')2
	Anti-Guinea Pig IgG Texas Red™ Conjugated Antibody, Goat Fab2 Anti-Guinea Pig IgG Texas
	Red™ Conjugated Antibody
	Background: F(ab')2 Anti-Guinea Pig IgG Texas Red Antibody was generated by enzymatic
	cleavage and subsequent separation from the Fc fragment. Because of their smaller size,
	F(ab)2 fragments offer several advantages over intact antibodies for use in certain
	immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue
	samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments
	lack the Fc region and therefore do not bind Fc receptors which effectively lowers background
	staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry,
	immunohistochemistry or IHC and other immunoassays.

### **Application Details**

Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	FLISA Dilution: 1:10,000 - 1:50,000
	Flow Cytometry Dilution: 1:500 - 1:2,500
	IF Microscopy Dilution: 1:1,000 - 1:5,000
Comment:	Post Translational Modification: Phosphorylation.
	Texas Red™ is a registered trademark of Molecular Probes Inc.
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

### Handling

	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing.  Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C prior to opening. Aliquot contents and freeze at -20 °C or below for extended storage. This product is stable for several weeks at 0 °C as an undiluted liquid.
Expiry Date:	12 months