

Datasheet for ABIN965063

Rabbit anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (Biotin) - Preadsorbed



Overview

Overview	
Quantity:	500 μg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Immunogen:	Immunogen: Guinea Pig IgG whole molecule
	Immunogen: Guinea Pig IgG whole molecule IgG
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Fragment:	IgG F(ab')2 fragment
Immunogen: Isotype: Fragment:	IgG F(ab')2 fragment Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat
Immunogen: Isotype: Fragment: Specificity:	IgG F(ab')2 fragment Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum.
Immunogen: Isotype: Fragment: Specificity: Purification:	IgG F(ab')2 fragment Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum.

Target Details

Target Type:	Antibody
Background:	Synonyms: Rabbit F(ab')2 Anti-Guinea Pig IgG Antibody Biotin Conjugation, Rabbit F(ab')2 Anti-
	Guinea Pig IgG Biotin Conjugated Antibody, Rabbit Fab2 Anti-Guinea Pig IgG Biotin Conjugated
	Antibody
	Background: F(ab')2 Anti-Guinea Pig IgG Biotin Antibody was generated by enzymatic cleavage
	and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2
	fragments offer several advantages over intact antibodies for use in certain immunochemical
	techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and
	show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc
	region and therefore do not bind Fc receptors which effectively lowers background staining.
	F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry,
	immunohistochemistry or IHC and other immunoassays.

Application Details

Application Notes:	Immunohistochemistry Dilution: 1:1,000 - 1:5,000
	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	ELISA Dilution: 1:90,000
	Western Blot Dilution: 1:2,000 - 1:10,000
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 500 µL Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	0.5 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

	should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing.
	This vial contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute
	1:10 by adding 225 μL of the buffer stated above directly to the vial. Recap, mix thoroughly and
	briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution
	when calculating final dilutions as recommended below.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening. Store the vial at -20 °C or below after dilution.
Expiry Date:	12 months