

Datasheet for ABIN965212

## Goat anti-Rabbit IgG (Heavy & Light Chain) Antibody - Preadsorbed

[Go to Product page](#)[1 Image](#)[2 Publications](#)

### Overview

Quantity:	500 µg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

### Product Details

Immunogen:	Immunogen: Rabbit IgG whole molecule
Isotype:	IgG
Fragment:	F(ab') <sub>2</sub> fragment
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabbit Serum.
Purification:	Preadsorption: Solid phase absorption
Sterility:	Sterile filtered

### Target Details

Target:	IgG
Abstract:	<a href="#">IgG Products</a>

## Target Details

---

Target Type:	Antibody
Background:	<p>Synonyms: Goat F(ab')<sub>2</sub> Anti-Rabbit IgG Antibody Pre-Adsorbed, Goat Fab<sub>2</sub> Anti-Rabbit IgG Antibody</p> <p>Background: F(ab')<sub>2</sub> Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)<sub>2</sub> fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)<sub>2</sub> fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)<sub>2</sub> fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')<sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.</p>

## Application Details

---

Application Notes:	<p>Immunohistochemistry Dilution: 1:1,000 - 1:5,000</p> <p>Application Note: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10<sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.</p> <p>ELISA Dilution: 1:20,000</p> <p>Western Blot Dilution: 1:2,000 - 1:10,000</p>
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Format:	Liquid
Concentration:	1.0 mg/mL
Buffer:	<p>Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</p> <p>Stabilizer: None</p> <p>Preservative: 0.01 % (w/v) Sodium Azide</p>
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

## Handling

should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Store vial at 4 °C prior to opening. This product is stable for several weeks at 4 °C as an undiluted liquid. For extended storage aliquot contents and freeze at -24 °C or below.


Expiry Date: 12 months

## Publications

Product cited in: Li, He, Zhang, Liu, Liu: "High expression of RAB38 promotes malignant progression of pancreatic cancer." in: **Molecular medicine reports**, Vol. 19, Issue 2, pp. 909-918, (2019) ([PubMed](#)).

Li, Wang, Li: "Kinesin family member 20B regulates tongue cancer progression by promoting cell proliferation." in: **Molecular medicine reports**, Vol. 19, Issue 3, pp. 2202-2210, (2019) ([PubMed](#)).

## Images



### SDS-PAGE

**Image 1.** SDS-PAGE results of Goat F(ab')<sub>2</sub> Anti-Rabbit IgG (H&L) Antibody. Lane 1: reduced F(ab')<sub>2</sub> anti-Mouse. Lane 2: reduced F(ab')<sub>2</sub> anti-Rabbit. Lane 3: Opal Prestained Ladder - MB-210-0500. Lane 4: non-reduced F(ab')<sub>2</sub> anti-Mouse. Lane 5: non-reduced F(ab')<sub>2</sub> anti-Rabbit. Load: 1.0ug. 4-20% Lonza SDS-PAGE; Coomassie Stained; BioRad ChemiDoc Imaged.