

Datasheet for ABIN965325

Goat anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (HRP)[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	HRP
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

Product Details

Purpose:	Fab Guinea Pig IgG (H&L) Antibody Peroxidase Conjugated
Immunogen:	Optional[Immunogen]: Guinea Pig IgG whole molecule
Isotype:	IgG
Fragment:	Fab fragment
Cross-Reactivity (Details):	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).
Characteristics:	Fab Anti-Guinea Pig IgG Fluorescein Antibody generated in goat detects guinea pig IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type,

Product Details

fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Target Details

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

Background: Fab Anti-Guinea Pig IgG Peroxidase Antibody generated in goat detects guinea pig IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Application Details

Application Notes: Application Note: Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Immunohistochemistry Dilution: 1:500 - 1:2,500 Western Blot Dilution: 1:1,000 - 1:5,000 ELISA Dilution: 1:10,000 - 1:50,000 Other: User Optimized

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution Buffer: Restore with deionized water (or equivalent), Reconstitution Volume: 1.0 mL

Concentration: 1.0 mg/mL

Buffer: Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
, Preservative: 0.01 % (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Preservative: Gentamicin sulfate

Handling

Precaution of Use:	This product contains Gentamicin sulfate: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months