

Datasheet for ABIN965327 Goat anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (Texas Red (TR)) - Preadsorbed



Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Texas Red (TR)
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)

Product Details

Immunogen:	Immunogen: Guinea Pig IgG whole molecule
lsotype:	lgG
Fragment:	Fab fragment
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum.
Characteristics:	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Purification:	Preadsorption: Solid phase absorption
Labeling Ratio:	3.1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN965327 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	lgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Goat Fab Anti-Guinea Pig IgG Texas Red™ Conjugated Antibody, Goat Fab Fragmer
	Anti-Guinea Pig IgG Antibody Texas Red™ Conjugation
	Background: Fab Anti-Guinea Pig IgG Texas Red Antibody generated in goat detects guinea pig
	IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both
	heavy and light chains of the antibody molecule are present. Secondary Antibodies are available
	in a variety of formats and conjugate types. When choosing a secondary antibody product,
	consideration must be given to species and immunoglobulin specificity, conjugate type,
	fragment and chain specificity, level of cross-reactivity, and host-species source and fragment
	composition.
Application Details	
Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibod
	based fluorescent assays requiring extremely low background levels, absence of F(c) mediated
	binding, lot-to-lot consistency, high titer and specificity.
	FLISA Dilution: 1:10,000 - 1:50,000
	Flow Cytometry Dilution: 1:500 - 1:2,500
	IF Microscopy Dilution: 1:1,000 - 1:5,000
Comment:	Texas Red™ is a registered trademark of Molecular Probes Inc.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL
	Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN965327 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Handling

	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening. Store the vial at -20 °C or below after dilution.
Expiry Date:	12 months