

## Datasheet for ABIN965333

## Goat anti-Golden Syrian Hamster IgG (Heavy & Light Chain) Antibody (TRITC) - Preadsorbed

1 mg



Go to Product page

	er		

Quantity:

quartity.	Ting
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Golden Syrian Hamster
Host:	Goat
Clonality:	Polyclonal
Conjugate:	TRITC
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)
Product Details	
Immunogen:	Immunogen: Golden Syrian Hamster IgG whole molecule
Isotype:	IgG
Fragment:	Fab fragment
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and
	anti-Goat Serum.
Characteristics:	anti-Goat Serum.  This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Characteristics:  Purification:	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex
	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

## **Target Details**

Concentration:

•			
Target:	IgG		
Abstract:	IgG Products		
Target Type:	Antibody		
Background:	Synonyms: Goat Fab Anti-Golden Syrian Hamster IgG Antibody Rhodamine Conjugation, Goat		
	Fab Anti-Golden Syrian Hamster IgG TRITC Conjugated Antibody, Goat Fab Anti-Hamster IgG		
	Antibody Rhodamine Conjugation, Goat Fab Anti-Hamster IgG TRITC Conjugated Antibody		
	Background: Fab Anti-Golden Syrian Hamster IgG (H&L) Antibody generated in goat detects		
	hamster IgG. Representing approximately 75 % of serum immunoglobulins, IgG is the most		
	abundant antibody isotype found in the circulation. IgG molecules are synthesized and secrete		
	by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate		
	types. When choosing a secondary antibody product, consideration must be given to species		
	and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-		
	reactivity, and host-species source and fragment composition. Fab Antibody is ideal for		
	investigators who routinely perform flow cytometry, immunofluorescence, IHC, and other		
	immunoassays. This Fab Anti-Golden Syrian Hamster IgG Antibody is conjugated to		
	Rhodamine.		
Application Details			
Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence		
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for		
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms		
	requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot		
	consistency, high titer and specificity.		
	FLISA Dilution: 1:10,000 - 1:50,000		
	Flow Cytometry Dilution: 1:500 - 1:2,500		
	IF Microscopy Dilution: 1:1,000 - 1:5,000		
Restrictions:	For Research Use only		
Handling			
Format:	Lyophilized		
Reconstitution:	Reconstitution Volume: 1.0 mL		

1.0 mg/mL

Reconstitution Buffer: Restore with deionized water (or equivalent)

## Handling

Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	
	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free	
	Preservative: 0.01 % (w/v) Sodium Azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Avoid cycles of freezing and thawing.	
	Product is photosensitive and should be protected from light.	
Storage:	RT,4 °C,-20 °C	
Storage Comment:	Store antibody at 4 °C prior to restoration. For extended storage aliquot contents and freeze at -	
	24 °C or below. This product is stable for several weeks at 4 °C as an undiluted liquid.	
Expiry Date:	12 months	