

## Datasheet for ABIN965383

# Goat anti-Rat IgG (Heavy & Light Chain) Antibody (Texas Red (TR)) - Preadsorbed



Go to Product page

Overview	

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Texas Red (TR)
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)
Draduat Dataila	
Product Details	
Immunogen:	Immunogen: Rat IgG whole molecule
	Immunogen: Rat IgG whole molecule  IgG
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Fragment:	IgG Fab fragment
Immunogen: Isotype: Fragment: Specificity:	IgG  Fab fragment  Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum.  This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex

#### **Target Details**

Target:	lgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Goat Fab Anti-Rat IgG Texas Red™ Conjugated Antibody, Goat Fab Fragment Anti-
	Rat IgG Antibody Texas Red™ Conjugation
	Background: Fab Anti-Rat IgG (H&L) Antibody generated in goat detects rat IgG. Representing
	approximately 75 % of serum immunoglobulins, IgG is the most abundant antibody isotype
	found in the circulation. IgG molecules are synthesized and secreted by plasma B cells.
	Secondary Antibodies are available in a variety of formats and conjugate types. When choosing
	a secondary antibody product, consideration must be given to species and immunoglobulin
	specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-
	species source and fragment composition. Fab Antibody is ideal for investigators who routinely
	perform flow cytometry, immunofluorescence, IHC, and other immunoassays. This Fab Anti-Rat
	IgG Antibody is conjugated to Texas Red™.

### **Application Details**

App	lication	Notes:
, (PP	iioa tioi i	I VOLCO.

Application Note: This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

FLISA Dilution: 1:10,000 - 1:50,000

Flow Cytometry Dilution: 1:500 - 1:2,500 IF Microscopy Dilution: 1:1,000 - 1:5,000

Comment:

Texas Red<sup>™</sup> is a registered trademark of Molecular Probes Inc.

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL
	Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL

## Handling

Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. Dilute only prior to immediate use This vial contains a relatively low volume of reagent (25 $\mu$ L). To minimize loss of volume dilute 1:10 by adding 225 $\mu$ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20 °C or below after dilution. Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening.
Expiry Date:	12 months