

Datasheet for ABIN965392

Rabbit anti-Sheep IgG (Heavy & Light Chain) Antibody (HRP)



(١,	er	٦/	iΔ	۱۸۱
_	ノV	\sim 1	٧		V V

Quantity:	500 μg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	HRP
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Purpose:	Fab Sheep IgG (H&L) Antibody Peroxidase Conjugated
Purpose: Immunogen:	Fab Sheep IgG (H&L) Antibody Peroxidase Conjugated Optional[Immunogen]: Sheep IgG whole molecule
Immunogen:	Optional[Immunogen]: Sheep IgG whole molecule
Immunogen:	Optional[Immunogen]: Sheep IgG whole molecule IgG

specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Fab Antibody is ideal for investigators who routinely perform flow cytometry, immunofluorescence, IHC, and other immunoassays. This Fab Antisheep IgG Antibody is conjugated to fluorescein.

Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Fab Anti-Sheep IgG (H&L) Antibody generated in rabbit detects immunoglobulin g from sheep,
	both heavy and light chains of the antibody molecule are present. Each IgG has two antigen
	binding sites. Representing approximately 75 % of serum immunoglobulins, IgG is the most
	abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted
	by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate
	types. When choosing a secondary antibody product, consideration must be given to species
	and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-
	reactivity, and host-species source and fragment composition. This Fab Anti-Sheep IgG
	Antibody is conjugated to Peroxidase.

Application Details

Application Notes:	Application Note: Fab Anti-Sheep IgG Peroxidase Antibody has been tested by ELISA and is

suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Immunohistochemistry Dilution: 1:500 - 1:2,500 Western Blot Dilution: 1:1,000 - 1:10,000 ELISA Dilution: 1:10,000 - 1:50,000 Other: User Optimized

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Buffer: Restore with deionized water (or equivalent), Reconstitution Volume: 500
	μL

Handling

Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free , Preservative: 0.01 % (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Preservative:	Gentamicin sulfate
Precaution of Use:	This product contains Gentamicin sulfate: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months