

Datasheet for ABIN965393

Rabbit anti-Sheep IgG (Heavy & Light Chain) Antibody (Biotin) - Preadsorbed



Go to Product page

Overview	
Quantity:	500 μg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Immunogen:	Immunogen: Sheep IgG whole molecule
Isotype:	IgG
Fragment:	Fab fragment
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin and anti-Rabbit Serum.
Purification:	Preadsorption: Solid phase absorption
Target Details	
Target:	IgG
Abstract:	IgG Products

Target Details

Target Type:	Antibody
Background:	Synonyms: Rabbit Fab Anti-Sheep IgG Biotin Conjugated Antibody, Rabbit Fab Fragment Anti-
	Sheep IgG Antibody Biotin Conjugation
	Background: Fab Anti-Sheep IgG (H&L) Antibody generated in rabbit detects immunoglobulin g
	from sheep, both heavy and light chains of the antibody molecule are present. Each IgG has
	two antigen binding sites. Representing approximately 75 $\%$ of serum immunoglobulins, IgG is
	the most abundant antibody isotype found in the circulation. IgG molecules are synthesized
	and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and
	conjugate types. When choosing a secondary antibody product, consideration must be given to
	species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of
	cross-reactivity, and host-species source and fragment composition. This Fab Anti-Sheep IgG
	Antibody is conjugated to Biotin.

Application Details

A 1: +! N - +	the control of the second case the second case.	DIL 1: 1.1 000 1.F 000
Application Notes:	immunonistochemistry	y Dilution: 1:1,000 - 1:5,000

Application Note: Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

ELISA Dilution: 1:20,000 - 1:100,000

Western Blot Dilution: 1:2,000 - 1:10,000

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 500 μ L Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	0.5 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. This vial contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening. Store the vial at -20 °C or below after dilution.
Expiry Date:	12 months